

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

1781



JAPANESE PATENT OFFICE

PATENT ABSTRACTS OF JAPAN

(11)Publication number: **10313896 A**

(43)Date of publication of
application: **02. 12 . 98**

(51)Int. Cl **C12Q 1/37**
// C07K 5/103
C07K 5/11
C07K 7/06

(21)Application number: 09126087

(22)Date of filing: 15 05 97

(71)Applicant: **YAMANOUCHI PHARMACEUT
CO LTD**

(72)Inventor: **KANO HIROYUKI**
SHIMIZU YASUAKI
KATO MASAO
SAITO SHIGEKI
MATSUMOTO SHUNICHIRO
AIBE KAZUHIKO

(54)NEW SYNTHETIC SUBSTRATE FOR
MEASUREMENT OF ACTIVITY HAVING
CHROMOPHORE OF FLUOROPHORE TO
MATRIX METALLOPROTEASE


(57)Abstract:

PROBLEM TO BE SOLVED: To obtain a new synthetic substrate necessary for screening of a matrix metalloprotease (MMP) inhibitor and suitable for rapid, simple and highly sensitive MMP assay system, etc., capable of carrying out multi-treatment by including a specific chemical structure.

SOLUTION: This compound is represented by formula I [Z is a protective group of N terminal; Y is a chromophore or a fluorophore; B is a peptide residue containing one or more amino acid residues hardly ingested with aminopeptidase; C is a peptide residue comprising an amino acid residue other than the amino acid residues hardly ingested with aminopeptidase], e.g.

Suc-Lys-Pro-Leu-Gly-Leu- Phe-Ala-Arg-MCA (MCA is 4-methyl-coumarin-7-yl-amide). Furthermore, activity of MMP and MMP containing double digesting process by aminopeptidase are preferably measured by using the synthetic substrate. The synthetic substrate is obtained by reacting, e.g. a compound of formula II with a compound of formula III in the presence of a condensing agent and then deprotecting the resultant reaction product.

COPYRIGHT: (C)1998,JPO



Ac-Lys(Boc)-Pro-Leu-Gly

II

Leu-Phe-Ala-Arg(pmc)-Glu

III